

COUNTY PROFILE: Boyd County, Kentucky

US COUNTY PERFORMANCE

The Institute for Health Metrics and Evaluation (IHME) at the University of Washington analyzed the performance of all 3,142 US counties or county-equivalents in terms of life expectancy at birth, mortality rates for select causes, alcohol use, smoking prevalence, obesity prevalence, and recommended physical activity using novel small area estimation techniques and the most up-to-date county-level information.

Explore more results using the interactive US Health Map data visualization (<http://vizhub.healthdata.org/subnational/usa>).

FINDINGS: LIFE EXPECTANCY

Sex	Boyd County	Kentucky	National	National rank	% change 1980-2014
Female	78.0	78.8	81.5	2627	+2.5
Male	73.9	73.7	76.7	2249	+6.9

life expectancy at birth (years), 2014

Fig. 1: Female life expectancy, 2014





Fig. 2: Male life expectancy, 2014



FINDINGS: ALL-CAUSE MORTALITY

Sex	Boyd County	Kentucky	National	National rank	% change 1980-2014
Female	881.4	827.2	667.8	2730	-7.2
Male	1182.5	1168.6	930.1	2463	-26.2

rate per 100,000 population, age-standardized, 2014

Fig. 3: Female all-cause mortality, 2014





Fig. 4: Male all-cause mortality, 2014



FINDINGS: ISCHEMIC HEART DISEASE

Sex	Boyd County	Kentucky	National	National rank	% change 1980-2014
Female	159.1	149.9	124.9	2276	-50.5
Male	271.2	235.3	191.5	2659	-53.0

rate per 100,000 population, age-standardized, 2014

Fig. 5: Female ischemic heart disease, 2014





Fig. 6: Male ischemic heart disease, 2014



FINDINGS: CEREBROVASCULAR DISEASE (STROKE)

Sex	Boyd County	Kentucky	National	National rank	% change 1980-2014
Female	76.9	55.3	47.4	3042	-11.5
Male	67.0	55.4	48.8	2752	-42.3

rate per 100,000 population, age-standardized, 2014

Fig. 7: Female cerebrovascular disease (stroke), 2014





Fig. 8: Male cerebrovascular disease (stroke), 2014



FINDINGS: TRACHEAL, BRONCHUS, AND LUNG CANCER

Sex	Boyd County	Kentucky	National	National rank	% change 1980-2014
Female	65.5	67.8	43.8	2907	+47.1
Male	100.5	113.1	67.6	2471	-24.6

rate per 100,000 population, age-standardized, 2014

Fig. 9: Female tracheal, bronchus, and lung cancer, 2014





Fig. 10: Male tracheal, bronchus, and lung cancer, 2014



FINDINGS: BREAST CANCER

Sex	Boyd County	Kentucky	National	National rank	% change 1980-2014
Female	32.5	27.6	25.9	2932	-14.0
Male	0.4	0.3	0.3	2428	-14.4

rate per 100,000 population, age-standardized, 2014

Fig. 11: Female breast cancer, 2014





Fig. 12: Male breast cancer, 2014



FINDINGS: MALIGNANT SKIN MELANOMA

Sex	Boyd County	Kentucky	National	National rank	% change 1980-2014
Female	2.3	2.3	1.9	2157	+2.2
Male	5.6	5.6	4.5	2379	+40.8

rate per 100,000 population, age-standardized, 2014

Fig. 13: Female malignant skin melanoma, 2014





Fig. 14: Male malignant skin melanoma, 2014



FINDINGS: DIABETES, UROGENITAL, BLOOD, AND ENDOCRINE DISEASES MORTALITY

Sex	Boyd County	Kentucky	National	National rank	% change 1980-2014
Female	53.4	62.9	49.6	1427	+18.8
Male	67.9	78.4	63.8	1612	+3.2

rate per 100,000 population, age-standardized, 2014

Fig. 15: Female diabetes, urogenital, blood, and endocrine diseases mortality, 2014





Fig. 16: Male diabetes, urogenital, blood, and endocrine diseases mortality, 2014



FINDINGS: SELF-HARM AND INTERPERSONAL VIOLENCE MORTALITY

Sex	Boyd County	Kentucky	National	National rank	% change 1980-2014
Female	8.7	10.3	9.0	1215	+2.2
Male	25.3	35.4	30.9	639	+3.3

rate per 100,000 population, age-standardized, 2014

Fig. 17: Female self-harm and interpersonal violence mortality, 2014





Fig. 18: Male self-harm and interpersonal violence mortality, 2014



FINDINGS: TRANSPORT INJURIES MORTALITY

Sex	Boyd County	Kentucky	National	National rank	% change 1980-2014
Female	13.6	12.7	8.1	1617	+9.6
Male	21.5	29.1	19.8	681	-30.2

rate per 100,000 population, age-standardized, 2014

Fig. 19: Female transport injuries mortality, 2014





Fig. 20: Male transport injuries mortality, 2014



FINDINGS: MENTAL AND SUBSTANCE USE DISORDERS MORTALITY

Sex	Boyd County	Kentucky	National	National rank	% change 1980-2014
Female	21.8	15.0	8.2	3062	+1842.1
Male	28.9	29.1	18.7	2840	+540.7

rate per 100,000 population, age-standardized, 2014

Fig. 21: Female mental and substance use disorders mortality, 2014





Fig. 22: Male mental and substance use disorders mortality, 2014



FINDINGS: CIRRHOSIS AND OTHER CHRONIC LIVER DISEASES MORTALITY

Sex	Boyd County	Kentucky	National	National rank	% change 1980-2014
Female	18.0	13.2	11.8	2816	+73.7
Male	27.3	25.4	22.2	2310	+16.4

rate per 100,000 population, age-standardized, 2014

Fig. 23: Female cirrhosis and other chronic liver diseases mortality, 2014





Fig. 24: Male cirrhosis and other chronic liver diseases mortality, 2014



FINDINGS: HEAVY DRINKING

Sex	Boyd County	Kentucky	National	National rank	% change 2005-2012
Female	4.1	4.6	6.7	1029	+65.4
Male	8.0	10.0	9.9	582	+56.6

prevalence (%), age-standardized, 2012

Fig. 25: Female heavy drinking, 2012





Fig. 26: Male heavy drinking, 2012



FINDINGS: BINGE DRINKING

Sex	Boyd County	Kentucky	National	National rank	% change 2002-2012
Female	8.1	9.5	12.4	860	+136.3
Male	19.2	21.0	24.5	578	+49.0

prevalence (%), age-standardized, 2012

Fig. 27: Female binge drinking, 2012





Fig. 28: Male binge drinking, 2012



FINDINGS: SMOKING

Sex	Boyd County	Kentucky	National	National rank	% change 1996-2012
Female	28.9	26.0	17.9	2986	+4.4
Male	29.9	28.8	22.2	2658	-15.6

prevalence (%), age-standardized, 2012

Fig. 29: Female smoking, 2012





Fig. 30: Male smoking, 2012



FINDINGS: OBESITY

Sex	Boyd County	Kentucky	National	National rank	% change 2001-2011
Female	40.0	39.3	36.1	1930	+24.8
Male	38.4	37.5	33.8	2028	+22.6

prevalence (%), age-standardized, 2011

Fig. 31: Female obesity, 2011





Fig. 32: Male obesity, 2011



FINDINGS: RECOMMENDED PHYSICAL ACTIVITY

Sex	Boyd County	Kentucky	National	National rank	% change 2001-2011
Female	43.9	45.8	52.6	2456	+33.2
Male	48.4	49.9	56.3	2464	+5.9

prevalence (%), age-standardized, 2011

Fig. 33: Female recommended physical activity, 2011





Fig. 34: Male recommended physical activity, 2011



CITATION:

Institute for Health Metrics and Evaluation (IHME),
US County Profile: Boyd County, Kentucky.
Seattle, WA: IHME, 2016.

Institute for Health Metrics and Evaluation
2301 Fifth Ave., Suite 600
Seattle, WA 98121 USA

Telephone: +1-206-897-2800
E-mail: engage@healthdata.org
www.healthdata.org

