
US COUNTY PERFORMANCE

The Institute for Health Metrics and Evaluation (IHME) at the University of Washington analyzed the performance of all 3,142 US counties or county-equivalents in terms of life expectancy at birth, mortality rates for select causes, alcohol use, smoking prevalence, obesity prevalence, and recommended physical activity using novel small area estimation techniques and the most up-to-date county-level information.

Explore more results using the interactive US Health Map data visualization (http://vizhub.healthdata.org/subnational/usa).


FINDINGS: LIFE EXPECTANCY

Sex	Breathitt County	Kentucky	National	National rank	% change 1980-2014
Female	73.7	78.8	81.5	3138	-3.7
Male	67.0	73.7	76.7	3135	+0.5



life expectancy at birth (years), 2014

Fig. 2: Male life expectancy, 2014



FINDINGS: ALL-CAUSE MORTALITY

Sex	Breathitt County	Kentucky	National	National rank	% change 1980-2014
Female	1215.4	827.2	667.8	3139	+40.6
Male	1709.1	1168.6	930.1	3136	+5.3

rate per 100,000 population, age-standardized, 2014

Fig. 3: Female all-cause mortality, 2014





Fig. 4: Male all-cause mortality, 2014



FINDINGS: ISCHEMIC HEART DISEASE

Sex	Breathitt County	Kentucky	National	National rank	% change 1980-2014
Female	329.3	149.9	124.9	3139	+8.3
Male	438.8	235.3	191.5	3134	-31.2

rate per 100,000 population, age-standardized, 2014




Fig. 6: Male ischemic heart disease, 2014 50 100 150 200 250 зóо 350 400 450 50⁰

FINDINGS: CEREBROVASCULAR DISEASE (STROKE)


Sex	Breathitt County	Kentucky	National	National rank	% change 1980-2014
Female	63.0	55.3	47.4	2611	-17.9
Male	66.8	55.4	48.8	2741	-32.0

rate per 100,000 population, age-standardized, 2014

40 50 60 70 80 90 100 110 120 130 20 30

Fig. 7: Female cerebrovascular disease (stroke), 2014

Fig. 8: Male cerebrovascular disease (stroke), 2014






FINDINGS: TRACHEAL, BRONCHUS, AND LUNG CANCER

Sex	Breathitt County	Kentucky	National	National rank	% change 1980-2014
Female	107.7	67.8	43.8	3135	+159.7
Male	189.1	113.1	67.6	3139	+52.1

rate per 100,000 population, age-standardized, 2014

Fig. 9: Female tracheal, bronchus, and lung cancer, 2014


Fig. 10: Male tracheal, bronchus, and lung cancer, 2014




FINDINGS: BREAST CANCER

Sex	Breathitt County	Kentucky	National	National rank	% change 1980-2014
Female	31.8	27.6	25.9	2856	+9.1
Male	0.4	0.3	0.3	2400	+6.6

rate per 100,000 population, age-standardized, 2014









Fig. 12: Male breast cancer, 2014


FINDINGS: MALIGNANT SKIN MELANOMA

Sex	Breathitt County	Kentucky	National	National rank	% change 1980-2014
Female	2.6	2.3	1.9	2817	+23.0
Male	6.0	5.6	4.5	2694	+72.0

rate per 100,000 population, age-standardized, 2014

Fig. 13: Female malignant skin melanoma, 2014





FINDINGS: DIABETES, UROGENITAL, BLOOD, AND ENDOCRINE DISEASES MORTALITY

Sex	Breathitt County	Kentucky	National	National rank	% change 1980-2014
Female	68.4	62.9	49.6	2466	+44.4
Male	73.7	78.4	63.8	1992	+51.4

rate per 100,000 population, age-standardized, 2014

Fig. 15: Female diabetes, urogenital, blood, and endocrine diseases mortality, 2014





Fig. 16: Male diabetes, urogenital, blood, and endocrine diseases mortality, 2014




FINDINGS: SELF-HARM AND INTERPERSONAL VIOLENCE MORTALITY

Sex	Breathitt County	Kentucky	National	National rank	% change 1980-2014
Female	14.4	10.3	9.0	2863	+33.0
Male	46.6	35.4	30.9	2782	-20.5

rate per 100,000 population, age-standardized, 2014

Fig. 17: Female self-harm and interpersonal violence mortality, 2014



FINDINGS: TRANSPORT INJURIES MORTALITY

Sex	Breathitt County	Kentucky	National	National rank	% change 1980-2014
Female	31.7	12.7	8.1	3121	+57.9
Male	85.4	29.1	19.8	3131	+38.3

rate per 100,000 population, age-standardized, 2014








Fig. 19: Female transport injuries mortality, 2014









FINDINGS: MENTAL AND SUBSTANCE USE DISORDERS MORTALITY

Sex	Breathitt County	Kentucky	National	National rank	% change 1980-2014
Female	32.0	15.0	8.2	3134	+2661.8
Male	63.4	29.1	18.7	3129	+1055.8

rate per 100,000 population, age-standardized, 2014

Fig. 21: Female mental and substance use disorders mortality, 2014





Fig. 22: Male mental and substance use disorders mortality, 2014



FINDINGS: CIRRHOSIS AND OTHER CHRONIC LIVER DISEASES MORTALITY

Sex	Breathitt County	Kentucky	National	National rank	% change 1980-2014
Female	13.7	13.2	11.8	1999	+81.0
Male	34.2	25.4	22.2	2839	+49.8

rate per 100,000 population, age-standardized, 2014

Fig. 23: Female cirrhosis and other chronic liver diseases mortality,





Fig. 24: Male cirrhosis and other chronic liver diseases mortality, 2014



FINDINGS: HEAVY DRINKING

Sex	Breathitt County	Kentucky	National	National rank	% change 2005-2012
Female	1.8	4.6	6.7	104	+88.4
Male	9.0	10.0	9.9	1043	+48.2

prevalence (%), age-standardized, 2012

Fig. 25: Female heavy drinking, 2012





Fig. 26: Male heavy drinking, 2012



FINDINGS: BINGE DRINKING

Sex	Breathitt County	Kentucky	National	National rank	% change 2002-2012
Female	5.3	9.5	12.4	180	+151.0
Male	17.4	21.0	24.5	306	+80.9

prevalence (%), age-standardized, 2012





Fig. 28: Male binge drinking, 2012




FINDINGS: SMOKING

Sex	Breathitt County	Kentucky	National	National rank	% change 1996-2012
Female	29.2	26.0	17.9	3002	-11.1
Male	35.8	28.8	22.2	3127	-0.4

prevalence (%), age-standardized, 2012

Fig. 29: Female smoking, 2012



FINDINGS: OBESITY

Sex	Breathitt County	Kentucky	National	National rank	% change 2001-2011
Female	50.3	39.3	36.1	3030	+33.8
Male	44.2	37.5	33.8	3106	+38.2

prevalence (%), age-standardized, 2011





Fig. 32: Male obesity, 2011



FINDINGS: RECOMMENDED PHYSICAL ACTIVITY

Sex	Breathitt County	Kentucky	National	National rank	% change 2001-2011
Female	37.8	45.8	52.6	2959	+29.8
Male	36.6	49.9	56.3	3131	-6.8

prevalence (%), age-standardized, 2011

Fig. 33: Female recommended physical activity, 2011





Fig. 34: Male recommended physical activity, 2011



CITATION:

Institute for Health Metrics and Evaluation (IHME), US County Profile: Breathitt County, Kentucky. Seattle, WA: IHME, 2016.

Institute for Health Metrics and Evaluation 2301 Fifth Ave., Suite 600 Seattle, WA 98121 USA Telephone: +1-206-897-2800 E-mail: engage@healthdata.org www.healthdata.org

