
# US COUNTY PERFORMANCE

The Institute for Health Metrics and Evaluation (IHME) at the University of Washington analyzed the performance of all 3,142 US counties or county-equivalents in terms of life expectancy at birth, mortality rates for select causes, alcohol use, smoking prevalence, obesity prevalence, and recommended physical activity using novel small area estimation techniques and the most up-to-date county-level information.

Explore more results using the interactive US Health Map data visualization (http://vizhub.healthdata.org/subnational/usa).


# FINDINGS: LIFE EXPECTANCY

| Sex    | Quay County | New Mexico | National | National rank | % change 1980-2014 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 79.4        | 81.1       | 81.5     | 2051          | +1.7               |
| Male   | 72.9        | 75.6       | 76.7     | 2552          | +6.5               |



life expectancy at birth (years), 2014

Fig. 2: Male life expectancy, 2014



# FINDINGS: ALL-CAUSE MORTALITY

| Sex    | Quay County | New Mexico | National | National rank | % change 1980-2014 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 789.0       | 663.0      | 667.8    | 2120          | -1.9               |
| Male   | 1169.4      | 952.2      | 930.1    | 2414          | -21.3              |

rate per 100,000 population, age-standardized, 2014





Fig. 4: Male all-cause mortality, 2014



# FINDINGS: ISCHEMIC HEART DISEASE

| Sex    | Quay County | New Mexico | National | National rank | % change 1980-2014 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 174.7       | 109.2      | 124.9    | 2566          | -33.0              |
| Male   | 234.8       | 172.3      | 191.5    | 2142          | -52.8              |

rate per 100,000 population, age-standardized, 2014





Fig. 6: Male ischemic heart disease, 2014



## FINDINGS: CEREBROVASCULAR DISEASE (STROKE)

| Sex    | Quay County | New Mexico | National | National rank | % change 1980-2014 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 41.4        | 42.1       | 47.4     | 468           | -40.8              |
| Male   | 43.7        | 39.6       | 48.8     | 636           | -50.3              |

rate per 100,000 population, age-standardized, 2014






Fig. 8: Male cerebrovascular disease (stroke), 2014





# FINDINGS: TRACHEAL, BRONCHUS, AND LUNG CANCER

| Sex    | Quay County | New Mexico | National | National rank | % change 1980-2014 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 41.3        | 30.8       | 43.8     | 950           | +37.0              |
| Male   | 70.2        | 45.4       | 67.6     | 1216          | -25.9              |

rate per 100,000 population, age-standardized, 2014






Fig. 10: Male tracheal, bronchus, and lung cancer, 2014




## FINDINGS: BREAST CANCER


| Sex    | Quay County | New Mexico | National | National rank | % change 1980-2014 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 25.8        | 23.3       | 25.9     | 1573          | -18.3              |
| Male   | 0.3         | 0.2        | 0.3      | 744           | -14.9              |

rate per 100,000 population, age-standardized, 2014



#### Fig. 11: Female breast cancer, 2014






# FINDINGS: MALIGNANT SKIN MELANOMA

| Sex    | Quay County | New Mexico | National | National rank | % change 1980-2014 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 2.0         | 1.8        | 1.9      | 1269          | -5.6               |
| Male   | 4.7         | 4.0        | 4.5      | 1393          | +23.6              |

rate per 100,000 population, age-standardized, 2014

Fig. 13: Female malignant skin melanoma, 2014



# FINDINGS: DIABETES, UROGENITAL, BLOOD, AND ENDOCRINE DISEASES MORTALITY

| Sex    | Quay County | New Mexico | National | National rank | % change 1980-2014 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 55.6        | 54.3       | 49.6     | 1596          | +56.0              |
| Male   | 77.5        | 70.2       | 63.8     | 2219          | +62.7              |

rate per 100,000 population, age-standardized, 2014

Fig. 15: Female diabetes, urogenital, blood, and endocrine diseases mortality, 2014





Fig. 16: Male diabetes, urogenital, blood, and endocrine diseases mortality, 2014





## FINDINGS: SELF-HARM AND INTERPERSONAL VIOLENCE MORTALITY


| Sex    | Quay County | New Mexico | National | National rank | % change 1980-2014 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 16.3        | 13.8       | 9.0      | 3010          | +25.9              |
| Male   | 50.1        | 46.5       | 30.9     | 2918          | -2.5               |

rate per 100,000 population, age-standardized, 2014

Fig. 17: Female self-harm and interpersonal violence mortality, 2014







# FINDINGS: TRANSPORT INJURIES MORTALITY

| Sex    | Quay County | New Mexico | National | National rank | % change 1980-2014 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 16.8        | 12.2       | 8.1      | 2248          | -24.4              |
| Male   | 41.1        | 29.1       | 19.8     | 2417          | -42.5              |

rate per 100,000 population, age-standardized, 2014





Fig. 20: Male transport injuries mortality, 2014



## FINDINGS: MENTAL AND SUBSTANCE USE DISORDERS MORTALITY

| Sex    | Quay County | New Mexico | National | National rank | % change 1980-2014 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 13.6        | 15.6       | 8.2      | 2695          | +465.6             |
| Male   | 44.2        | 38.9       | 18.7     | 3082          | +169.9             |

rate per 100,000 population, age-standardized, 2014

Fig. 21: Female mental and substance use disorders mortality, 2014





Fig. 22: Male mental and substance use disorders mortality, 2014




## FINDINGS: CIRRHOSIS AND OTHER CHRONIC LIVER DISEASES MORTALITY


| Sex    | Quay County | New Mexico | National | National rank | % change 1980-2014 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 23.8        | 23.0       | 11.8     | 3066          | +78.0              |
| Male   | 55.2        | 41.3       | 22.2     | 3107          | +60.5              |

rate per 100,000 population, age-standardized, 2014

Fig. 23: Female cirrhosis and other chronic liver diseases mortality,



#### Fig. 24: Male cirrhosis and other chronic liver diseases mortality, 2014



## FINDINGS: HEAVY DRINKING

| Sex    | Quay County | New Mexico | National | National rank | % change 2005-2012 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 2.8         | 5.8        | 6.7      | 439           | +46.6              |
| Male   | 6.8         | 8.3        | 9.9      | 187           | -0.7               |

Fig. 25: Female heavy drinking, 2012

prevalence (%), age-standardized, 2012




Fig. 26: Male heavy drinking, 2012

## FINDINGS: BINGE DRINKING

| Sex    | Quay County | New Mexico | National | National rank | % change 2002-2012 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 6.8         | 10.3       | 12.4     | 525           | +0.2               |
| Male   | 18.6        | 22.4       | 24.5     | 486           | -23.1              |

prevalence (%), age-standardized, 2012




Fig. 27: Female binge drinking, 2012





Fig. 28: Male binge drinking, 2012

## FINDINGS: SMOKING

| Sex    | Quay County | New Mexico | National | National rank | % change 1996-2012 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 19.6        | 18.9       | 17.9     | 938           | -14.1              |
| Male   | 25.8        | 23.6       | 22.2     | 1666          | -7.9               |

Fig. 29: Female smoking, 2012 20 10 15 25 зо 35 40


prevalence (%), age-standardized, 2012



FINDINGS: OBESITY

| Sex    | Quay County | New Mexico | National | National rank | % change 2001-2011 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 40.3        | 34.9       | 36.1     | 1986          | +17.3              |
| Male   | 37.0        | 31.2       | 33.8     | 1373          | +35.8              |

prevalence (%), age-standardized, 2011






Fig. 32: Male obesity, 2011

## FINDINGS: RECOMMENDED PHYSICAL ACTIVITY

| Sex    | Quay County | New Mexico | National | National rank | % change 2001-2011 |
|--------|-------------|------------|----------|---------------|--------------------|
| Female | 48.5        | 54.5       | 52.6     | 1853          | +8.6               |
| Male   | 52.6        | 57.8       | 56.3     | 1756          | +2.4               |

prevalence (%), age-standardized, 2011





Fig. 34: Male recommended physical activity, 2011



## **CITATION:**

Institute for Health Metrics and Evaluation (IHME), US County Profile: Quay County, New Mexico. Seattle, WA: IHME, 2016.

Institute for Health Metrics and Evaluation 2301 Fifth Ave., Suite 600 Seattle, WA 98121 USA Telephone: +1-206-897-2800 E-mail: engage@healthdata.org www.healthdata.org

