
## US COUNTY PERFORMANCE

The Institute for Health Metrics and Evaluation (IHME) at the University of Washington analyzed the performance of all 3,142 US counties or county-equivalents in terms of life expectancy at birth, mortality rates for select causes, alcohol use, smoking prevalence, obesity prevalence, and recommended physical activity using novel small area estimation techniques and the most up-to-date county-level information.

Explore more results using the interactive US Health Map data visualization (http://vizhub.healthdata.org/subnational/usa).


#### FINDINGS: LIFE EXPECTANCY

Sex	Swisher County	Texas	National	National rank	% change 1980-2014
Female	80.6	80.8	81.5	1450	+2.6
Male	75.5	76.2	76.7	1616	+7.2



life expectancy at birth (years), 2014

Fig. 2: Male life expectancy, 2014



## FINDINGS: ALL-CAUSE MORTALITY

Sex	Swisher County	Texas	National	National rank	% change 1980-2014
Female	733.0	694.2	667.8	1611	-8.4
Male	1001.4	954.8	930.1	1498	-27.7

rate per 100,000 population, age-standardized, 2014

Fig. 3: Female all-cause mortality, 2014





Fig. 4: Male all-cause mortality, 2014



# FINDINGS: ISCHEMIC HEART DISEASE

Sex	Swisher County	Texas	National	National rank	% change 1980-2014
Female	126.0	127.2	124.9	1356	-50.2
Male	188.9	197.7	191.5	1123	-63.2

rate per 100,000 population, age-standardized, 2014

Fig. 5: Female ischemic heart disease, 2014 . 50 100 150 200 250 зóо 350




Fig. 6: Male ischemic heart disease, 2014

## FINDINGS: CEREBROVASCULAR DISEASE (STROKE)

Sex	Swisher County	Texas	National	National rank	% change 1980-2014
Female	51.2	52.0	47.4	1558	-54.1
Male	66.2	54.0	48.8	2719	-46.3

rate per 100,000 population, age-standardized, 2014





Fig. 8: Male cerebrovascular disease (stroke), 2014



# FINDINGS: TRACHEAL, BRONCHUS, AND LUNG CANCER

Sex	Swisher County	Texas	National	National rank	% change 1980-2014
Female	31.5	38.4	43.8	265	+40.1
Male	53.4	62.9	67.6	425	-36.5

rate per 100,000 population, age-standardized, 2014





Fig. 9: Female tracheal, bronchus, and lung cancer, 2014 Fig. 10: Male tra



# Fig. 10: Male tracheal, bronchus, and lung cancer, 2014

#### FINDINGS: BREAST CANCER

Sex	Swisher County	Texas	National	National rank	% change 1980-2014
Female	25.3	25.6	25.9	1382	-7.9
Male	0.3	0.3	0.3	364	-14.9

rate per 100,000 population, age-standardized, 2014




Fig. 11: Female breast cancer, 2014





Fig. 12: Male breast cancer, 2014

# FINDINGS: MALIGNANT SKIN MELANOMA

Sex	Swisher County	Texas	National	National rank	% change 1980-2014
Female	1.9	1.6	1.9	952	-6.9
Male	4.4	4.0	4.5	994	+23.0

rate per 100,000 population, age-standardized, 2014



C 5 8 2 ່ສ 4 ż ģ

# FINDINGS: DIABETES, UROGENITAL, BLOOD, AND ENDOCRINE DISEASES MORTALITY

Sex	Swisher County	Texas	National	National rank	% change 1980-2014
Female	64.7	55.9	49.6	2260	+83.5
Male	93.6	66.9	63.8	2829	+111.6

rate per 100,000 population, age-standardized, 2014

Fig. 15: Female diabetes, urogenital, blood, and endocrine diseases mortality, 2014




Fig. 16: Male diabetes, urogenital, blood, and endocrine diseases mortality, 2014





Fig. 14: Male malignant skin melanoma, 2014

#### FINDINGS: SELF-HARM AND INTERPERSONAL VIOLENCE MORTALITY

Sex	Swisher County	Texas	National	National rank	% change 1980-2014
Female	8.3	8.4	9.0	1042	-15.0
Male	28.4	28.8	30.9	1010	-29.8

rate per 100,000 population, age-standardized, 2014

Fig. 17: Female self-harm and interpersonal violence mortality, 2014



FINDINGS: TRANSPORT INJURIES MORTALITY

Sex	Swisher County	Texas	National	National rank	% change 1980-2014
Female	15.1	9.5	8.1	1922	-11.3
Male	35.6	24.3	19.8	2013	-30.1

rate per 100,000 population, age-standardized, 2014





Fig. 20: Male transport injuries mortality, 2014

Fig. 18: Male self-harm and interpersonal violence mortality, 2014





20

40

60

80

100

120

140

160

180

#### FINDINGS: MENTAL AND SUBSTANCE USE DISORDERS MORTALITY

Sex	Swisher County	Texas	National	National rank	% change 1980-2014
Female	6.3	6.4	8.2	1040	+356.7
Male	7.7	13.9	18.7	242	+115.1

rate per 100,000 population, age-standardized, 2014





Fig. 21: Female mental and substance use disorders mortality, 2014 Fig. 22: Male mental and substance use disorders mortality, 2014




#### FINDINGS: CIRRHOSIS AND OTHER CHRONIC LIVER DISEASES MORTALITY


Sex	Swisher County	Texas	National	National rank	% change 1980-2014
Female	16.0	14.5	11.8	2547	+88.5
Male	29.0	27.7	22.2	2495	+57.6

rate per 100,000 population, age-standardized, 2014

Fig. 23: Female cirrhosis and other chronic liver diseases mortality,



#### Fig. 24: Male cirrhosis and other chronic liver diseases mortality, 2014



## FINDINGS: HEAVY DRINKING

Sex	Swisher County	Texas	National	National rank	% change 2005-2012
Female	2.6	5.3	6.7	349	+34.7
Male	6.2	9.5	9.9	94	-1.5

Fig. 25: Female heavy drinking, 2012

prevalence (%), age-standardized, 2012




Fig. 26: Male heavy drinking, 2012

#### FINDINGS: BINGE DRINKING

Sex	Swisher County	Texas	National	National rank	% change 2002-2012
Female	7.7	10.8	12.4	759	+18.5
Male	20.5	24.5	24.5	845	-0.6

prevalence (%), age-standardized, 2012






Fig. 28: Male binge drinking, 2012




## FINDINGS: SMOKING

Sex	Swisher County	Texas	National	National rank	% change 1996-2012
Female	17.2	15.1	17.9	425	-13.4
Male	23.9	22.1	22.2	1099	-13.8

Fig. 29: Female smoking, 2012 10 15 20 25 зо 35 40






FINDINGS: OBESITY

Sex	Swisher County	Texas	National	National rank	% change 2001-2011
Female	40.8	38.9	36.1	2092	+20.0
Male	37.8	36.0	33.8	1754	+23.3

prevalence (%), age-standardized, 2011






Fig. 32: Male obesity, 2011

#### FINDINGS: RECOMMENDED PHYSICAL ACTIVITY

Sex	Swisher County	Texas	National	National rank	% change 2001-2011
Female	45.4	49.5	52.6	2276	+10.5
Male	49.9	54.1	56.3	2234	+3.0

prevalence (%), age-standardized, 2011





Fig. 34: Male recommended physical activity, 2011



#### **CITATION:**

Institute for Health Metrics and Evaluation (IHME), US County Profile: Swisher County, Texas. Seattle, WA: IHME, 2016.

Institute for Health Metrics and Evaluation 2301 Fifth Ave., Suite 600 Seattle, WA 98121 USA Telephone: +1-206-897-2800 E-mail: engage@healthdata.org www.healthdata.org

